

3.2 - Forest growth/dynamics (theory)

Miquel De Cáceres, Victor Granda, Aitor Ameztegui

Ecosystem Modelling Facility

2022-06-15



Outline


1. Simulation models and processes
2. Carbon pools and carbon balance
3. Growth and senescence
4. Mortality and recruitment

1. Simulation models and processes

Models in medfate and their processes

Simulation functions in **medfate** have been designed and implemented in a **nested** manner:

- Each upper-level function includes the processes of lower-level ones and adds new processes.
- Consequently, the two water balance models (basic and advanced) are available for simulations with `growth()` and `fordyn()`.
- In this presentation, we will focus on the state variables and processes that are added in simulations with `growth()` and `fordyn()`.



1. Simulation models and processes

The `growth()` and `fordyn()` models

The `growth()` model implements several **daily** processes:

- Carbon balance processes that modulate changes in *leaf area*, *sapwood area* and *fine root area* of woody plant cohorts.

1. Simulation models and processes

The `growth()` and `fordyn()` models

The `growth()` model implements several **daily** processes:

- Carbon balance processes that modulate changes in *leaf area*, *sapwood area* and *fine root area* of woody plant cohorts.
- Structural changes (diameter and height for trees; cover and height for shrubs) derived from growth and senescence processes.

1. Simulation models and processes

The `growth()` and `fordyn()` models

The `growth()` model implements several **daily** processes:

- Carbon balance processes that modulate changes in *leaf area*, *sapwood area* and *fine root area* of woody plant cohorts.
- Structural changes (diameter and height for trees; cover and height for shrubs) derived from growth and senescence processes.
- Tree mortality, which reduces cohort density.

1. Simulation models and processes

The `growth()` and `fordyn()` models

The `growth()` model implements several **daily** processes:

- Carbon balance processes that modulate changes in *leaf area*, *sapwood area* and *fine root area* of woody plant cohorts.
- Structural changes (diameter and height for trees; cover and height for shrubs) derived from growth and senescence processes.
- Tree mortality, which reduces cohort density.

For applications spanning more than a few years, users will use the `growth()` model indirectly, via calls to `fordyn()`, which includes other processes implemented at **annual** steps:

- Natural regeneration (recruitment)

1. Simulation models and processes

The `growth()` and `fordyn()` models

The `growth()` model implements several **daily** processes:

- Carbon balance processes that modulate changes in *leaf area*, *sapwood area* and *fine root area* of woody plant cohorts.
- Structural changes (diameter and height for trees; cover and height for shrubs) derived from growth and senescence processes.
- Tree mortality, which reduces cohort density.

For applications spanning more than a few years, users will use the `growth()` model indirectly, via calls to `fordyn()`, which includes other processes implemented at **annual** steps:

- Natural regeneration (recruitment)
- The effect of disturbances (at present, only forest management)

2. Carbon compartments, pools and carbon balance

Carbon compartments and pools

Types

Structural carbon - Cell membranes, cell walls & cytosolic machinery.

Metabolic carbon - Labile carbon (sugar) concentration used to sustain cell functioning.

Storage carbon - Labile carbon (starch) concentration used as long-term carbon source.

2. Carbon compartments, pools and carbon balance

Carbon compartments and pools

Types

Structural carbon - Cell membranes, cell walls & cytosolic machinery.

Metabolic carbon - Labile carbon (sugar) concentration used to sustain cell functioning.

Storage carbon - Labile carbon (starch) concentration used as long-term carbon source.

Pools

Compartment	Structural	Metabolic	Storage
Leaves	Leaf dry biomass B_{leaf}	Leaf sugar SS_{leaf}	Leaf starch ST_{leaf}
Sapwood (branch/stem/coarse roots)	Sapwood dry biomass $B_{sapwood}$	Sapwood sugar $SS_{sapwood}$	Sapwood starch $ST_{sapwood}$
Fine roots	Fine root dry biomass $B_{fineroot}$		

2. Carbon compartments, pools and carbon balance

Carbon compartments and pools

Types

Structural carbon - Cell membranes, cell walls & cytosolic machinery.


Metabolic carbon - Labile carbon (sugar) concentration used to sustain cell functioning.

Storage carbon - Labile carbon (starch) concentration used as long-term carbon source.

Pools

Compartment	Structural	Metabolic	Storage
Leaves	Leaf dry biomass B_{leaf}	Leaf sugar SS_{leaf}	Leaf starch ST_{leaf}
Sapwood (branch/stem/coarse roots)	Sapwood dry biomass $B_{sapwood}$	Sapwood sugar $SS_{sapwood}$	Sapwood starch $ST_{sapwood}$
Fine roots	Fine root dry biomass $B_{fineroot}$		

Fluxes



2. Carbon compartments, pools and carbon balance

Labile carbon balance equations

A. Changes in **leaf metabolic** carbon result from considering gross photosynthesis (A_g), leaf maintenance respiration (M_{leaf}), phloem transport (F_{phloem}) and sugar-starch dynamics (SC_{leaf}) and translocation to sapwood storage (TS_{leaf}):

$$\Delta SS_{leaf} \cdot V_{storage,leaf} = A_g - M_{leaf} - F_{phloem} - SC_{leaf} - TS_{leaf}$$

2. Carbon compartments, pools and carbon balance

Labile carbon balance equations

A. Changes in **leaf metabolic** carbon result from considering gross photosynthesis (A_g), leaf maintenance respiration (M_{leaf}), phloem transport (F_{phloem}) and sugar-starch dynamics (SC_{leaf}) and translocation to sapwood storage (TS_{leaf}):

$$\Delta SS_{leaf} \cdot V_{storage,leaf} = A_g - M_{leaf} - F_{phloem} - SC_{leaf} - TS_{leaf}$$

B. Changes in **leaf storage** carbon result from sugar-starch dynamics and translocation to sapwood storage (TT_{leaf}):

$$\Delta ST_{leaf} \cdot V_{storage,leaf} = SC_{leaf} - TT_{leaf}$$

2. Carbon compartments, pools and carbon balance

Labile carbon balance equations

A. Changes in **leaf metabolic** carbon result from considering gross photosynthesis (A_g), leaf maintenance respiration (M_{leaf}), phloem transport (F_{phloem}) and sugar-starch dynamics (SC_{leaf}) and translocation to sapwood storage (TS_{leaf}):

$$\Delta SS_{leaf} \cdot V_{storage,leaf} = A_g - M_{leaf} - F_{phloem} - SC_{leaf} - TS_{leaf}$$

B. Changes in **leaf storage** carbon result from sugar-starch dynamics and translocation to sapwood storage (TT_{leaf}):

$$\Delta ST_{leaf} \cdot V_{storage,leaf} = SC_{leaf} - TT_{leaf}$$

C. Changes in **stem metabolic** carbon result from considering phloem transport, maintenance respiration of sapwood ($M_{sapwood}$) and fineroot ($M_{fineroot}$) tissues, sugar-starch dynamics and translocation to sapwood storage ($TS_{sapwood}$):

$$\Delta SS_{sapwood} \cdot V_{storage,sapwood} = F_{phloem} - M_{sapwood} - M_{fineroot} - SC_{sapwood} - TS_{sapwood}$$

2. Carbon compartments, pools and carbon balance

Labile carbon balance equations

A. Changes in **leaf metabolic** carbon result from considering gross photosynthesis (A_g), leaf maintenance respiration (M_{leaf}), phloem transport (F_{phloem}) and sugar-starch dynamics (SC_{leaf}) and translocation to sapwood storage (TS_{leaf}):

$$\Delta SS_{leaf} \cdot V_{storage,leaf} = A_g - M_{leaf} - F_{phloem} - SC_{leaf} - TS_{leaf}$$

B. Changes in **leaf storage** carbon result from sugar-starch dynamics and translocation to sapwood storage (TT_{leaf}):

$$\Delta ST_{leaf} \cdot V_{storage,leaf} = SC_{leaf} - TT_{leaf}$$

C. Changes in **stem metabolic** carbon result from considering phloem transport, maintenance respiration of sapwood ($M_{sapwood}$) and fineroot ($M_{fineroot}$) tissues, sugar-starch dynamics and translocation to sapwood storage ($TS_{sapwood}$):

$$\Delta SS_{sapwood} \cdot V_{storage,sapwood} = F_{phloem} - M_{sapwood} - M_{fineroot} - SC_{sapwood} - TS_{sapwood}$$

C. Changes in **stem storage** carbon result from considering sugar-starch dynamics, translocation from other pools, growth respiration and root exudation:


$$\Delta ST_{sapwood} \cdot V_{storage,sapwood} = SC_{sapwood} + TS_{leaf} + TT_{leaf} + TS_{sapwood} - G_{sapwood} - G_{leaf} - G_{fineroot} - RE_{sapwood}$$

2. Carbon compartments, pools and carbon balance

Labile carbon balance equations

Changes in labile carbon pools can be reduced to the balance between gross photosynthesis (A_g), maintenance (M), growth (G) and root exudation (RE) components:

$$\Delta S_{labile} = A_g - M - G - RE$$



2. Carbon compartments, pools and carbon balance

Design of labile carbon processes

- **Phloem transport** is only explicitly modelled when using the **advanced water submodel**, following Hölttä et al. (2017).

2. Carbon compartments, pools and carbon balance

Design of labile carbon processes

- **Phloem transport** is only explicitly modelled when using the **advanced water submodel**, following Hölttä et al. (2017).
- **Sugar/starch dynamics** are meant to keep sugar levels at a prescribed equilibrium value.

2. Carbon compartments, pools and carbon balance

Design of labile carbon processes

- **Phloem transport** is only explicitly modelled when using the **advanced water submodel**, following Hölttä et al. (2017).
- **Sugar/starch dynamics** are meant to keep sugar levels at a prescribed equilibrium value.
- The usual separation between **maintenance** and **growth** respiration is followed, with maintenance costs depending on temperature and phenology.

2. Carbon compartments, pools and carbon balance

Design of labile carbon processes

- **Phloem transport** is only explicitly modelled when using the **advanced water submodel**, following Hölttä et al. (2017).
- **Sugar/starch dynamics** are meant to keep sugar levels at a prescribed equilibrium value.
- The usual separation between **maintenance** and **growth** respiration is followed, with maintenance costs depending on temperature and phenology.
- **Translocation** of labile carbon towards sapwood storage occurs whenever there is senescence in a given tissue (leaves, branches).

2. Carbon compartments, pools and carbon balance

Design of labile carbon processes

- **Phloem transport** is only explicitly modelled when using the **advanced water submodel**, following Hölttä et al. (2017).
- **Sugar/starch dynamics** are meant to keep sugar levels at a prescribed equilibrium value.
- The usual separation between **maintenance** and **growth** respiration is followed, with maintenance costs depending on temperature and phenology.
- **Translocation** of labile carbon towards sapwood storage occurs whenever there is senescence in a given tissue (leaves, branches).
- **Root exudation** is not a process competing for metabolic carbon, but a consequence of plant storage capacity being surpassed (Prescott et al. 2020).

2. Carbon compartments, pools and carbon balance

Biomass balance

Balance in structural carbon pools

The change in structural biomass of each **compartment** results from the interplay between growth and senescence:

$$\Delta B_{leaves} = B_{leaves,growth} - B_{leaves,senescence}$$

$$\Delta B_{sapwood} = B_{sapwood,growth} - B_{sapwood,senescence}$$

$$\Delta B_{fineroot} = B_{fineroot,growth} - B_{fineroot,senescence}$$

2. Carbon compartments, pools and carbon balance

Biomass balance

Balance in structural carbon pools

The change in structural biomass of each **compartment** results from the interplay between growth and senescence:

$$\Delta B_{leaves} = B_{leaves,growth} - B_{leaves,senescence}$$

$$\Delta B_{sapwood} = B_{sapwood,growth} - B_{sapwood,senescence}$$

$$\Delta B_{fineroot} = B_{fineroot,growth} - B_{fineroot,senescence}$$

The following equation defines the structural biomass balance at the plant level:

$$\Delta B_{structure} = \Delta B_{leaves} + \Delta B_{sapwood} + \Delta B_{fineroot}$$

2. Carbon compartments, pools and carbon balance

Biomass balance

Balance in structural carbon pools

The change in structural biomass of each **compartment** results from the interplay between growth and senescence:

$$\Delta B_{leaves} = B_{leaves,growth} - B_{leaves,senescence}$$

$$\Delta B_{sapwood} = B_{sapwood,growth} - B_{sapwood,senescence}$$

$$\Delta B_{fineroot} = B_{fineroot,growth} - B_{fineroot,senescence}$$

The following equation defines the structural biomass balance at the plant level:

$$\Delta B_{structure} = \Delta B_{leaves} + \Delta B_{sapwood} + \Delta B_{fineroot}$$

Balance at the individual level

The biomass balance at the **individual level** is completed if we add the labile biomass balance to the structural biomass balance:

$$\Delta B_{plant} = \Delta B_{structure} + \Delta S_{labile}$$

2. Carbon compartments, pools and carbon balance

Biomass balance

Balance at the cohort level

At the **cohort level** we need to take into account that some individuals will die, so that the biomass balance needs to incorporate mortality losses:

$$\Delta B_{cohort} = \Delta B_{plant} \cdot N_{cohort} - B_{mortality}$$

where N_{cohort} is the initial cohort density (before mortality occurred) and $B_{mortality}$ is the biomass loss due to mortality of individuals, which in order to close the balance has to be defined as:

$$B_{mortality} = (B_{plant} + \Delta B_{plant}) \cdot N_{dead}$$

where N_{dead} is the density of dead individuals and B_{plant} is the initial plant biomass.

3. Growth and senescence

Leaf area/fine root biomass growth

Daily leaf area increment ΔLA and fine root biomass increment $\Delta B_{fineroot}$ are defined as the minimum of three constraints:

$$\Delta LA = \min(\Delta LA_{alloc}, \Delta LA_{source}, \Delta LA_{sink})$$

$$\Delta B_{fineroot} = \min(\Delta B_{fineroot,alloc}, \Delta B_{fineroot,source}, \Delta B_{fineroot,sink})$$

1. ΔLA_{alloc} and $\Delta B_{fineroot,alloc}$ are the increments allowed according to the targets set by **allocation rules**.
2. ΔLA_{source} and $\Delta B_{fineroot,source}$ are the maximum increments allowed by current **carbon availability**.
3. ΔLA_{sink} and $\Delta B_{fineroot,sink}$ are the increments expected by taking into account maximum growth rates as well as **temperature and turgor limitations** (Cabon et al. 2020).

3. Growth and senescence

Sapwood area growth

Daily sapwood formation, ΔSA , is defined as the minimum of two values expressing source and sink constraints:

$$\Delta SA = \min(\Delta SA_{source}, \Delta SA_{sink})$$

3. Growth and senescence

Sapwood area growth

Daily sapwood formation, ΔSA , is defined as the minimum of two values expressing source and sink constraints:

$$\Delta SA = \min(\Delta SA_{source}, \Delta SA_{sink})$$

Unlike leaf area or fine root biomass, sapwood formation is not explicitly constrained by any allocation rule.

3. Growth and senescence

Sapwood area growth

Daily sapwood formation, ΔSA , is defined as the minimum of two values expressing source and sink constraints:

$$\Delta SA = \min(\Delta SA_{source}, \Delta SA_{sink})$$

Unlike leaf area or fine root biomass, sapwood formation is not explicitly constrained by any allocation rule.

However, newly assimilated carbon is preferentially allocated to leaves and fine roots whenever storage levels are low because ΔSA_{source} is more restrictive than ΔLA_{source} or $\Delta B_{fineroot,source}$.

3. Growth and senescence

Key growth parameters

Parameter	R	Definition	Explanation
$RGR_{cambium,max}$	RGRcambiummax	Maximum daily tree sapwood growth rate relative to cambium perimeter length	Determines overall maximum growth rates for tree species
$RGR_{sapwood,max}$	RGRsapwoodmax	Maximum daily shrub sapwood growth rate relative to cambium perimeter length	Determines overall maximum growth rates for shrub species
$1/H_v$	Al2As	Leaf area to sapwood area ratio	Determines allocation target for leaves
RGR	Ar2Al	Root area to leaf area ratio	Determines allocation target for fine roots and influences root maintenance costs
$RSSG$	RSSG	Minimum relative starch for sapwood growth to occur	Determines preference for maintenance over growth under low carbon availability (e.g. shade-tolerant species)

3. Growth and senescence

Drivers of tissue senescence

Compartment	Senescence cause	Description	Key parameter
Leaf	Aging	Defoliation in winter-deciduous. Evergreens lose leaves progressively or suddenly, but with an amount related to leaf lifespan	LeafDuration
	Cavitation	Increases in stem PLC are translated to proportional leaf area losses	
Sapwood	Aging	Daily rate of conversion from sapwood to heartwood, depending on temperature and plant height	SRsapwood
Fine root	Aging	Daily rate of senescence depending on temperature	SRfineroot

3. Growth and senescence

Drivers of tissue senescence

Compartment	Senescence cause	Description	Key parameter
Leaf	Aging	Defoliation in winter-deciduous. Evergreens lose leaves progressively or suddenly, but with an amount related to leaf lifespan	LeafDuration
	Cavitation	Increases in stem PLC are translated to proportional leaf area losses	
Sapwood	Aging	Daily rate of conversion from sapwood to heartwood, depending on temperature and plant height	SRsapwood
Fine root	Aging	Daily rate of senescence depending on temperature	SRfineroot

Note: Stem cavitation reduces functional sapwood but does not lead to heartwood formation.

3. Growth and senescence

Updating structural variables

Trees

New sapwood area, ΔSA , is translated to an increment in DBH, ΔDBH , following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH$$

3. Growth and senescence

Updating structural variables

Trees

New sapwood area, ΔSA , is translated to an increment in DBH, ΔDBH , following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH$$

Increments in height are linearly related to increments in diameter through a coefficient depending on light conditions (Lindner et al. 1997), and are limited by a maximum height.

3. Growth and senescence

Updating structural variables

Trees

New sapwood area, ΔSA , is translated to an increment in DBH, ΔDBH , following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH$$

Increments in height are linearly related to increments in diameter through a coefficient depending on light conditions (Lindner et al. 1997), and are limited by a maximum height.

Crown ratios are updated following static allometric relationships

3. Growth and senescence

Updating structural variables

Trees

New sapwood area, ΔSA , is translated to an increment in DBH, ΔDBH , following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH$$

Increments in height are linearly related to increments in diameter through a coefficient depending on light conditions (Lindner et al. 1997), and are limited by a maximum height.

Crown ratios are updated following static allometric relationships

Shrubs

Leaf area changes are translated to changes in shrub volume, cover and shrub height via allometric equations.

3. Growth and senescence

Updating structural variables

Trees

New sapwood area, ΔSA , is translated to an increment in DBH, ΔDBH , following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH$$

Increments in height are linearly related to increments in diameter through a coefficient depending on light conditions (Lindner et al. 1997), and are limited by a maximum height.

Crown ratios are updated following static allometric relationships

Shrubs

Leaf area changes are translated to changes in shrub volume, cover and shrub height via allometric equations.

As for trees, shrub height is limited to a maximum value.

4. Mortality and recruitment

Plant mortality

Dynamic vegetation models implement mortality processes in very different ways, from purely-empirical to process-based (Hawkes 2000).

4. Mortality and recruitment

Plant mortality

Dynamic vegetation models implement mortality processes in very different ways, from purely-empirical to process-based (Hawkes 2000).

At present, plant mortality is formulated in an overly simplistic way and requires development efforts!

1. Species-specific parameters determine a daily constant **baseline mortality rate**.

4. Mortality and recruitment

Plant mortality

Dynamic vegetation models implement mortality processes in very different ways, from purely-empirical to process-based (Hawkes 2000).

At present, plant mortality is formulated in an overly simplistic way and requires development efforts!

1. Species-specific parameters determine a daily constant **baseline mortality rate**.
2. Daily mortality rates increase due to **carbon starvation** whenever sapwood metabolic carbon, $SS_{sapwood}$, becomes lower than a threshold, set by default to 30% of the homeostatic (equilibrium) concentrations.

4. Mortality and recruitment

Plant mortality

Dynamic vegetation models implement mortality processes in very different ways, from purely-empirical to process-based (Hawkes 2000).

At present, plant mortality is formulated in an overly simplistic way and requires development efforts!

1. Species-specific parameters determine a daily constant **baseline mortality rate**.
2. Daily mortality rates increase due to **carbon starvation** whenever sapwood metabolic carbon, $SS_{sapwood}$, becomes lower than a threshold, set by default to 30% of the homeostatic (equilibrium) concentrations.
3. Daily mortality rates increase due to **desiccation** whenever stem relative water content becomes lower than a pre-specified threshold, set to 30% by default.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).
2. **Mean temperature of the coldest month** is above a bioclimatic threshold.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).
2. **Mean temperature of the coldest month** is above a bioclimatic threshold.
3. The **fraction of light reaching the ground** is above a bioclimatic threshold.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).
2. **Mean temperature of the coldest month** is above a bioclimatic threshold.
3. The **fraction of light reaching the ground** is above a bioclimatic threshold.
4. The **moisture index** (annual precipitation over annual evapotranspiration) is larger than a bioclimatic threshold.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).
2. **Mean temperature of the coldest month** is above a bioclimatic threshold.
3. The **fraction of light reaching the ground** is above a bioclimatic threshold.
4. The **moisture index** (annual precipitation over annual evapotranspiration) is larger than a bioclimatic threshold.

When these conditions are met, a **probability of recruitment** is used to stochastically determine whether recruitment actually occurs.

4. Mortality and recruitment

Recruitment

Unlike mortality, recruitment is simulated *at the end of a simulated year* (only in `fordyn()`), following the design of many forest gap models.

The following conditions need to be met for a species to recruit:

1. Trees of the species are present with a **size above a maturity threshold** (or a seed rain is forced).
2. **Mean temperature of the coldest month** is above a bioclimatic threshold.
3. The **fraction of light reaching the ground** is above a bioclimatic threshold.
4. The **moisture index** (annual precipitation over annual evapotranspiration) is larger than a bioclimatic threshold.

When these conditions are met, a **probability of recruitment** is used to stochastically determine whether recruitment actually occurs.

Plants are recruited with specified structural characteristics (DBH and height) depending on species parameters.

M.C. Escher - Three worlds, 1955

